arXiv:0907.0701v1 [math.RT] 3 Jul 2009

Laura string algebras

Julie Dionne [a,1],

[a] *Département de mathématiques, Université de Sherbrooke, 2500, boul. de l'Université, Sherbrooke, Québec, Canada, J1K 2R1*

Abstract

We give a simple combinatorial criterion allowing to recognize whether a string (or, more generally, a special biserial) algebra is a laura algebra or not. We also show that a special biserial algebra is laura if and only if it has a finite number of isomorphism classes of indecomposable modules which have projective dimension and injective dimension greater than or equal to two, solving a conjecture ok Skowroński for special biserial algebras.

Key words: laura algebras, string algebras, special biserial algebras
MSC : 16S35, 18E30

Let k be an algebraically closed field. A finite-dimensional k-algebra R is biserial if the radical of every projective indecomposable R-module is the sum of two uniserial modules whose intersection is simple or zero [7]. In 1983, Skowroński and Waschbüsch characterized biserial algebras of finite representation type by the fact that almost split sequences have at most two non-projective middle terms [12]. The biserial algebras which have at most two middle terms in their almost split sequences are called string algebras (see [6]). The following definition is equivalent:

Definition 1 *A k-algebra R is a string algebra if it admits a presentation $R = kQ/I$ such that :*

(1) Each point has at most two arrows entering and two arrows exiting;
(2) For each arrow $\alpha : x \to y$ there is at most one arrow $\beta : y \to z$ such that $\alpha\beta$ is not in I and at most one arrow $\gamma : z \to x$ such that $\gamma\alpha$ is not in I;
(3) The ideal I is monomial.

Email address: `julie.dionne2@usherbrooke.ca` (Julie Dionne).

[1] Partially supported by Université de Sherbrooke.

The characterizations of string algebras which are of finite representation type [6], tilted [10], quasi-tilted [9] and shod [5] are known. For a string algebra $R = kQ/I$, we have the following definitions. A walk in the quiver of R is *reduced* if it contains no subwalk of the form $\alpha\alpha^{-1}$ or of the form $\alpha^{-1}\alpha$. A *cycle* is a non-oriented cycle, that is, a reduced walk starting and ending in the same point of the quiver. A walk ω is a *string* if it is reduced and contains no relation. Given a string ω, the representation admitting a copy of the field k at x for each passage of ω on x and with the obvious morphisms is the *string module* over ω, denoted by $M(\omega)$. A walk ω is a *band* if it is a cyclic string which is not the power of another cyclic string and if there is no n such that ω^n is in I.

Laura algebras have been defined independently by Assem and Coelho [2] and Skowroński [11]. Let A be an algebra, its left part is the full subcategory of $\mathrm{ind}A$ defined as follows:

$$\mathcal{L}_A = \{M \in \mathrm{ind}A \mid \text{ for all } \ L \rightsquigarrow M,\ \mathrm{dp}L \leq 1\}$$

Its right part is dually defined and is denoted by $\mathcal{R}_A$.

Definition 2 *An algebra A is laura if $\mathrm{ind}A\backslash(\mathcal{L}_A \cup \mathcal{R}_A)$ contains finitely many objects.*

We say that R is *strict laura* if it is laura but not quasi-tilted. Skowroński conjectured that an algebra R is laura if and only if the number of indecomposable modules with projective and injective dimension greater than or equal to two is finite.

Our aim is to characterize laura string algebras and to show Skowroński's conjecture for these algebras.

Definition 3 *Let $R \cong kQ/I$ be a string algebra, and ω be a double-zero on Q (see [9] for instance). We say that ω is an interlaced double-zero, abbreviated by DOZE, if there exist a band ω_2, two walks ω_1 and ω_3 and two relations ρ_1 and ρ_2 such that $\omega = \rho_1\omega_1\omega_2\omega_3\rho_2$.*

Remark that if $\omega = \rho_1\omega_1\omega_2\omega_3\rho_2$ is a DOZE, then $\rho_1\omega_1{\omega_2}^n\omega_3\rho_2$ is a double-zero for all $n \geq 0$.

The first two sections will be useful for proving the following theorem. Recall that an algebra is quasi-tilted of canonical type if its bounded derived category is equivalent to the bounded derived category of a category of coherent sheaves $\mathrm{coh}\mathbb{X}$ on a weighted projective line $\mathbb{X}$ in the sense of Geigle and Lenzing.

Theorem 4 *Let $R = kQ/I$ be a string algebra having no DOZE and such that Q contains a band.*

a) If Q has at least one band with at least one arrow entering and at least one arrow exiting, then R is quasi-tilted of canonical type.
b) If Q contains at least one band and is such that each band has either only exiting arrows, or only entering arrows, then R is strict laura or tilted.

In the third section, we prove the validity of Skowroński's conjecture for string and more generally for special biserial algebras. Our main results are the following two theorems.

Theorem 5 *Let $R = kQ/I$ be a string algebra. The following are equivalent:*

a) There is no DOZE on (Q, I);
b) $R = kQ/I$ is laura;
c) R has only a finite number of isomorphism classes of indecomposable modules having projective and injective dimension greater than or equal to 2.

If R is a special biserial algebra, let J be the minimal ideal containing all paths lying in non-monomial relation. Then R/J is a string algebra.

Theorem 6 *Let $R = kQ/I$ be a special biserial algebra. The following are equivalent:*

a) There is no DOZE on (Q, I);
b) $R = kQ/I$ is laura;
c) R has only a finite number of indecomposable modules having a projective and an injective dimension greater or equal than 2;
d) R/J is laura.

1 Preliminaries

A quiver Q is a tuple $(Q_0,\ Q_1,\ s : Q_1 \to Q_0,\ t : Q_1 \to Q_0)$. We call Q_0 the set of vertices, Q_1 the set of arrows, $s(\alpha)$ the source of the arrow α and $t(\alpha)$ the target of the arrow α. The algebra kQ is the k-vector space generated by all the paths on Q with the multiplication defined by the composition of paths. If I is an ideal of kQ, consider the algebra kQ/I. We have a complete set of idempotents of kQ/I given by the trivial paths on each vertex x, denoted by ε_x. For a finite dimensional, basic and connected k-algebra $R \cong kQ/I$, we denote by $\operatorname{mod} R$ the category of finite dimensional left R-modules. The indecomposable projective, injective and simple modules associated to ε_x will be denoted respectively by P_x, I_x and S_x. A path in $\operatorname{mod} R$ from M to N is a sequence

$$(*) M = M_0 \xrightarrow{f_1} M_1 \longrightarrow \cdots \qquad \xrightarrow{f_t} M_t = N$$

of non-zero morphisms between indecomposable modules. The path $(*)$ is called sectional if, for any i, we have that $X_i \not\cong \tau X_{i+2}$. A refinement of $(*)$ is a path

$$(*) M = X_0 \xrightarrow{g_1} X_1 \longrightarrow \cdots \qquad \xrightarrow{g_s} X_s = N$$

with $s \geq r$ such that there is an order-preserving function $\sigma : \{1, 2, 3, ..., t-1\} \to \{1, 2, 3, ..., s-1\}$ such that $M_i \cong X_{\sigma(i)}$ for all i. The path $(*)$ is called sectionally refinable if it has a sectional refinement. It is well known that an algebra is laura if and only if there is only a finite number of isomorphism classes of indecomposable modules on a path going from an injective module to a projective module. Let R be a strict laura or a tilted algebra then R has a unique faithful nonsemiregular component which is quasidirected. Let $({}_l\Gamma_\lambda)_{\lambda\in\Lambda}$ be the left stable parts of the faithful non semiregular components (since R is laura, Λ is a finite set). Let $({}_l\Sigma_\lambda)_{\lambda\in\Lambda}$ be complete slices of each one of those stable parts (since each component has a finite number of orbits, such slices exist). For all $\lambda \in \Lambda$, we define ${}_\infty R_\lambda$ to be the full subcategory generated by the support of ${}_l\Sigma_\lambda$. The left end algebra ${}_\infty R$ of R is by definition the product of the ${}_\infty R_\lambda$. We define dually the right end algebra R_∞ of R.

2 String algebras without DOZE

Throughout this section, let $R = kQ/I$ be a string algebra. The following lemma will be useful for the next two sections:

Lemma 7 *If $R = kQ/I$ has no DOZE, then two bands of R intersect in at most one point.*

Proof: We verify that for all cases with more than one point in common, we obtain a DOZE. If ω_1 and ω_2 have a non trivial walk u in common, we have :



or



where $\omega_1 = u\alpha_1\omega_1'\beta_1$ and $\omega_2 = u{\alpha_2}^{-1}\omega_2'{\beta_2}^{-1}$ in the first case and $\omega_1 = u\alpha_1\omega_1'{\beta_1}^{-1}$ and $\omega_2 = u{\alpha_2}^{-1}\omega_2'\beta_2$ in the second case. In the first case, $\alpha_2\omega_1'\beta_2$ is a DOZE. In the second, $\alpha_2\omega_1'u\omega_2'\beta_1$ is a DOZE. If ω_1 and ω_2 have no non trivial walks in common, we have one of the following three cases:

1)



2)



3)



where $\omega_1 = \beta_1 u\gamma_1\delta_1\omega'_1\alpha_1$ and $\omega_2 = \beta_2 v\gamma_2\delta_2\omega'_2\alpha_2$ in the first case, $\omega_1 = {\beta_1}^{-1}u\gamma_1\delta_1\omega'_1\alpha_1$ and $\omega_2 = \beta_2 v\gamma_2\delta_2\omega'_2{\alpha_2}^{-1}$ in the second case and $\omega_1 = {\beta_1}^{-1}u\gamma_1{\delta_1}^{-1}\omega'_1\alpha_1$ and $\omega_2 = \beta_2 v{\gamma_2}^{-1}\delta_2\omega'_2{\alpha_2}^{-1}$ in the last case.

Then, in the first case $\alpha_1\beta_1 u\gamma_1\delta_1$ is a DOZE, in the second $\beta_1\beta_2 v\gamma_2\delta_2$ is a DOZE, and in the third $\beta_1\beta_2 v{\gamma_2}^{-1}\delta_1^{-1}w\alpha_1\alpha_2$ is a DOZE.
□

2.1 Quasi-tilted algebras without DOZE

Remark 8 *Let $R = kQ/I$ be a string algebra without DOZE having a band Θ with entering arrow β and exiting arrow α. Let β^+ be an arrow of Θ such that $\beta\beta^+ \in I$. Then there is a string ω on Θ, going from the source of β^+ to the source of α, such that $\omega\alpha$ is a string.*

Proposition 9 *Let Q be a quiver with a band having at least one entering arrow β and at least one exiting arrow α. If $R = kQ/I$ is a string algebra having no DOZE, then R has no double-zero.*

Proof: We denote by β^+ the arrow such that $\beta\beta^+ \in I$ and by α^- the arrow such that $\alpha^-\alpha \in I$. Suppose that we have a double-zero. If $\beta\beta^+$ is the relation at the beginning of the double-zero then we have a DOZE. Thus, suppose that this is not the case. Then we can find a walk of minimal length between the end point of β^+ and a point on the double-zero. If the first relation on the walk of minimal length composed with a part of the double-zero points in the same direction than $\beta\beta^+$, we have a DOZE. Otherwise, let ω be the walk going from $t(\beta)$ to $s(\alpha^-)$ such that $\beta\omega$ is a string. Then we have a DOZE.
□

2.2 Cycles on a string algebra without DOZE

We now consider the case where each cycle has only entering arrows or only exiting arrows. We ignore the trivial case where R has only one band with neither entering nor exiting arrow (in this case, R is hereditary). We denote by $\Theta_1, \Theta_2, ..., \Theta_n$ the bands (up to a cyclic permutation) having only exiting arrows. Since R is finite dimensional, a band Θ has a point a such that all arrows starting at a belong to Θ. We arbitrarily choose such a point on each cycle Θ_i and denote it by a_i. Let α be an exiting arrow of Θ_i. We denote by α^- the arrow of Θ_i such that $\alpha^-\alpha \in I$. We denote by $\rho'_{(i,\alpha)}$ the minimal reduced walk going from a_i to $t(\alpha)$ and passing through $\alpha^-\alpha$.

We denote by $\Delta_1, \Delta_2, ..., \Delta_m$ the bands (up to a cyclic permutation) having only entering arrows.

Definition 10 *Let $R = kQ/I$ and ω be a string of (Q, I). We denote by $W(\omega)$ the set of strings ω' of (Q, I) such that $\omega' = \omega_1\omega\omega_2$. We denote by $D(\omega)$ the subcategory of (Q, I) whose objects are the points x such that there exists a string in $W(\omega)$ passing through x and whose morphisms are the composition of arrows α for which there exists a string of $W(\omega)$ passing through α.*

Definition 11 *Let $R = kQ/I$ be a string algebra such that each band on its quiver has only entering arrows or only exiting arrows. We define A_i to be the subcategory of R whose objects are the points of $D(\varepsilon_{a_i})$ and whose morphisms are given by the linear combinations of paths of arrows of $D(\varepsilon_{a_i})$. The algebra A_i is the quotient of the path algebra of $D(\varepsilon_{a_i})$ by $I \cap D(\varepsilon_{a_i})$. Note that A_i does not depend on the choice of a_i.*

We define dually the categories B_j.

Example: Let Q be the quiver



with I the ideal generated by $\alpha_1\rho_1$, $\alpha_2\rho_4$, $\rho_5\delta_1$, $\rho_8\delta_2$, $\beta_i\alpha_i$ for i such that $i \in \{1, 2\}$, $\gamma_i\beta_i$ for i such that $\in \{1, 2\}$ and $\delta_j\gamma_j = 0$ for j such that $j \in \{1, 2\}$. Then $R = Q/I$ contains no DOZE. We have $B = B_1 \times B_2$ with B_1 the full subcategory generated by $\{1, 2, 5\}$ and B_2 the full subcategory generated by $\{3, 4, 6\}$. We also have $A = A_1 \times A_2$ with A_1 the full subcategory generated by $\{8, 10, 11\}$ and A_2 the full subcategory generated by $\{9, 12, 13\}$.

Lemma 12 *The categories $D(\varepsilon_{a_i})$ are full in R.*

Proof: By contradiction, suppose that x and y are objects of A_i and let $\gamma : x \rightarrow y$ be an arrow which is not in $D(\varepsilon_{a_i})$. Then there exist strings $\omega : a_i \rightsquigarrow x$ and $\omega' : a_i \rightsquigarrow y$, and since γ is not in A_i, ω and ω' are such that $\omega\gamma$ and $\omega'\gamma^{-1}$ are reduced walks containing a zero-relation. But in this case, the zero-relation must contain γ and thus $\omega = u\alpha_1 ... \alpha_n$ and $\omega' = u'\alpha_m^{-1} ... \alpha_{n+2}^{-1}$, with $\alpha_1 ... \alpha_n\gamma$ and $\gamma\alpha_{n+2} ... \alpha_m$ some relations. Thus, $\gamma\omega'^{-1}\Theta_i\rho'_{(i,\alpha)}$ is a DOZE, a contradiction. We can apply the same proof to every morphism in the category R.
$\square$

Lemma 13 *Let* $x \in (Q_{A_i})_0$ *and* $y \notin (Q_{A_i})_0$. *Then there is no arrow* $\alpha : y \rightarrow x$. *As a consequence, if a relation* ρ *does not start in* $(Q_{A_i})_0$ *then it does not end in* $(Q_{A_i})_0$.

Proof: Suppose that such an arrow exists. Let ω be a string from a_i to x. Then $\omega\alpha^{-1}$ is not a reduced walk or else contains a zero-relation. If it is not a reduced walk, then $\omega = \omega'\alpha$, and $y \in (Q_{A_i})_0$, a contradiction. If it contains a zero-relation, then $\omega = \omega'\beta_n^{-1} ... \beta_1^{-1}$ with $\alpha\beta_1 ... \beta_n$ in I and so $\alpha\beta_1 ... \beta_n\omega'^{-1}\rho'_{i,\gamma}$, with γ an exiting arrow of Θ_i, is a DOZE, another contradiction.
$\square$

Corollary 14 *The categories* $D(\varepsilon_{a_i})$ *are convex in* R.

Proof: This follows directly from lemma 13.
$\square$

Lemma 15 *Each of the categories* $D(\varepsilon_{a_i})$ *contains only one simple cycle. In particular, it has no oriented cycle. Therefore, no relation* ρ *starts in* $(Q_{A_i})_0$ *and ends outside the cycle* Θ_i.

Proof: Suppose that there are two distinct cycles Θ and Θ_i in $D(\varepsilon_{a_i})$. If Θ contains a relation, we have a DOZE by gluing the relation with a walk relating the two cycles and which is chosen such that the two relations point in the same direction (this is possible because Θ and Θ_i are cycles). If not, we have a cycle with an exiting or an entering arrow and so we have a relation of the form $\alpha^{-}\alpha$ or of the form $\beta\beta^{+}$. In each case, we can construct a DOZE. The last statement follows from Lemma 13.
$\square$

Recall that an algebra R is left glued if $\mathcal{R}_R$ is cofinite in ind$R$ (see [1]).

Proposition 16 *For all* i, *with* $1 \leq i \leq n$, *the algebra* A_i *is a tilted algebra containing a complete slice in its postprojective component. In particular, it is left glued and all but a finite number of isomorphism classes of indecomposable* A_i*-modules are in* $\mathcal{R}_{A_i}$.

Proof: Let α be an arrow from x to y in A_i. We want to show that the projective A_i-modules P_x and P_y, with respective tops S_x and S_y, are in the same component of the Auslander-Reiten quiver of A_i. The radical of P_x is the direct sum of at most two terms, one of them is a uniserial module L with top S_y. Moreover, there is an epimorphism f from P_y to L. Since the inclusion of the radical of a projective module into this projective module is irreducible, we only have to show that f lies in a finite power of the radical of the module category.

Suppose $f_1 : M \to L$ and $f_2 : P_y \to M$ is a factorisation of f, where M may be decomposable.

$$\begin{tikzcd} P_y \arrow[r, "f_2"] & M \arrow[r, "f_1"] & L \arrow[r] & P_x . \end{tikzcd}$$

Since f is an epimorphism, L is a quotient of M. Let z_1 belong to the support of L, z_2 belong to the support of M but not to that of L and β be an arrow from z_1 to z_2. Since L is a direct summand of radP_x, there exists a walk ω' such that $\omega'\beta$ is in the support of M and $\alpha\omega'\beta \in I$ (ω' may be trivial).

We have a relation of I starting at x, thus x is on the cycle Θ_i by Lemma 15. Since there is no relation on a band, β is not on a cycle (by Lemma 15), and every arrow on a reduced walk beginning with $\alpha\omega'\beta$ and being the successor of β is not in Θ_i.

Thus, every arrow of the support of M lies in the support of L or is not in Θ_i. Since Θ_i is not included in the support of L, it is not in the support of M.

The k-dimensions of the direct summands of M are bounded above since the support of M contains no band.

We showed that there exist only a finite number of isomorphism classes of indecomposable modules which are direct summands of a module through which f factors.

Now, we only have to see that A_i contains no double-zero (otherwise we have a DOZE since we have a string between every point of A_i and Θ_i). So A_i is tilted (see [10]) and we have our result by Theorem 3.4 of [10].

□

Definition 17 *Let E be the set of strings which are not in one of the subcategories A_i or B_j. We define the middle part C to be the subcategory which has as set of objects $\bigcup_{\omega\in E}(D(\omega))_0$ and as set of morphisms $\Sigma_{\omega\in E}(D(\omega)(x,y))$ from x to y.*

Remark 18 *The algebra C is of finite representation type since it contains no band and it is a string algebra (see [6]).*

2.3 Module Category

We now begin the study of the category of indecomposable modules of R, with the aim to prove that it is a laura algebra under assumptions of this subsection. We denote by $A = \prod_{i=1}^{n} A_i$ the emphright side algebra of R and by $B = \prod_{j=1}^{m} B_j$ the emphleft side algebra of R.

Proposition 19 $\mathrm{ind}R = \mathrm{ind}A \cup \mathrm{ind}B \cup \mathrm{ind}C$

Proof: Every A_i-module or B_j-module can be considered as an R-module since A_i and B_j are full and convex subcategories of R, by completing their representations by zeros. Every C-module is a string module and thus is a string R-module.
Let M be an indecomposable R-module and ω be a string lying in the support of M. Then each point and each arrow of the support of M are in $D(\omega)$. Suppose that all points of M are in one of the A_i, or in one of the B_j, but not all in the same. Let x be a point of A_i which is not in B_j and y be a point of B_j which is not in A_i. Then the string $\omega : x \rightsquigarrow y$ cannot be in A_i, nor in B_j, and thus by definition lies in C. $\square$

Proposition 20 *Let M be an indecomposable A_i-module, N be an indecomposable R-module which is not an A_i-module and $f : M \to N$ be a non-zero morphism. Then M is a C-module or M is a B-module.*

Proof: Suppose that M is neither a B-module, nor a C-module. Then the support of M contains at least a point z' which is not an object of $B \cup C$. So it is an object of A_i for some i.
The image of f is a submodule of N, so there exist a string ω going from z' to x and an arrow $\alpha : y \to x$, with x in the support of the image of f and y in the support of N but not in the support of the image of f.
Since N is not an A-module, the support of N contains at least a point z which is not an object of A_i. Since z and y are in the support of N, there exists a string ν going from y to z of the form.

$$\begin{tikzcd}
z' \arrow[r, squiggly, no head, "\omega"] & x & y \arrow[l, "\alpha"'] \arrow[r, squiggly, no head, "\nu"] & z \, .
\end{tikzcd}$$

But z' is not in $B \cup C$ and z is not in A_i. Then there is no string between them, since such a string would be neither in A, nor in B, so it would be in C. If $\omega\alpha^{-1}\nu$ is a reduced walk, it contains a relation and this relation must contain α. But in this case, we obtain a relation entering in A_i, a contradiction.
Thus $\omega\alpha^{-1}\nu$ is not reduced and the relation between z and z' goes in the other direction. Suppose that $\alpha_1\alpha_2...\alpha_n$ is going from z'^* to z^*, as illustrated bellow.

$$\begin{tikzcd}
z'^* \arrow[r, "\alpha_1"] \arrow[rrrr, dotted, bend left, no head] & \cdots \arrow[r, "\alpha_i"] & \cdot \arrow[r, "\alpha_{i+1}"] \arrow[d, squiggly, no head, "\eta"] & \cdots \arrow[r, "\alpha_n"] & z^* \, . \\
 & & x & & \\
 & & y \arrow[u] & &
\end{tikzcd}$$

Since ω and ν are strings, so are $\alpha_1 ... \alpha_i \eta$ and $\eta^{-1}\alpha_{i+1}...\alpha_n$. Since R is a string algebra, this implies that $n = 2$.

$$\begin{tikzcd}
z'^{*} \arrow[r, "\alpha_1"] \arrow[rr, bend left, dotted, no head] & \cdot \arrow[r, "\alpha_2"] \arrow[d, squiggly, no head, "\eta"] & z^{*} . \\
 & x & \\
 & y \arrow[u] &
\end{tikzcd}$$

The target of α_1 is on ω and on ν, so it is in the support of Imf. Since Imf is a quotient of M, z'^{*} is in Imf. And since Imf is a submodule of N, z^{*} is in Imf, a contradiction to the fact that $\alpha_1\alpha_2 \in I$ and to Lemma 15.
□

Proof of Theorem 4: The first part of Theorem 4 follows now directly from [10] (Theorems 2.6 and 3.4) and [8] (Theorem 3.1).

For the second part, the last proposition with the fact that $|\text{ind}A_i \cap \text{ind}B_j| \leq |\text{ind}(A_i \cap B_j)| < \infty$ say that there exists only a finite number of isomorphism classes of indecomposable A_i-modules which admit successors from ind$R\backslash$ indA_i. Let $\mathcal{X}_i$ be the subcategory of indA_i which contains modules without successors from ind$R\backslash$ indA_i, that is

$$\mathcal{X}_i = \{X \in \textit{ind}A_i \ / \ \textit{for all } 0 \neq f : X \to Y \ \textit{in indR}, \ Y \in \ \textit{ind}A_i\}.$$

We have that $\mathcal{X}_i$ is such that if M is an A_i-module in $\mathcal{X}_i$ and if there is a non-zero morphism f from M to M', then M' is in $\mathcal{X}_i$. We define dually $\mathcal{Y}_j$.

We have that if $f : M \to N$ is a morphism, where M and N are objects of $\mathcal{X}_i$, then f is irreducible in A_i if and only if f is irreducible in R, since the morphisms are preserved (this follows from the fullness of the categories A_i and B_j and from the fact that there are no points lying in $A_i \cap A_j$ if $i \neq j$). By studying the Auslander-Reiten translation in A_i and in R, we can show that it is preserved. Take

$$(*) : \qquad \begin{tikzcd} 0 \arrow[r] & M \arrow[r] & {}_R I_0 \arrow[r, "i_1"] & {}_R I_1 \end{tikzcd}$$

a minimal injective resolution of M in R. We know that $\tau_R^{-1}(M) = \text{Coker } \nu^{-1}(i_1)$, where ν denotes the Nakayama functor.

$$\begin{tikzcd} \nu^{-1}({}_R I_0) \arrow[r, "\nu^{-1}(i_1)"] & \nu^{-1}({}_R I_1) \arrow[r] & \text{Coker } \nu^{-1}(i_1) \end{tikzcd}$$

We compute $\tau_{A_i}^{-1}(M)$ in the same way. In general, $\nu^{-1}({}_{A_i} I_0)$ is a quotient of $\nu^{-1}({}_R I_0)$ and $\nu^{-1}({}_{A_i} I_1)$ is a quotient of $\nu^{-1}({}_R I_1)$. The kernels K_0 and K_1 of those projections contain no points of A_i in their supports.

$$\begin{tikzcd}
0 \arrow[d] & 0 \arrow[d] & & \\
K_0 \arrow[r] \arrow[d] & K_1 \arrow[r] \arrow[d] & L \arrow[r] \arrow[d] & 0 \\
\nu^{-1}({}_R I_0) \arrow[r, "\nu^{-1}(i_1)"] \arrow[d] & \nu^{-1}({}_R I_1) \arrow[r] \arrow[d] & \tau_R^{-1}(M) \arrow[r] \arrow[d, "g"] & 0 \\
\nu^{-1}({}_{A_i} I_0) \arrow[r, "\nu^{-1}(i_1)"] \arrow[d] & \nu^{-1}({}_{A_i} I_1) \arrow[r] \arrow[d] & \tau_{A_i}^{-1}(M) \arrow[r] & 0 \\
0 & 0 & &
\end{tikzcd}$$

Let L be the cokernel of the induced morphism from K_0 to K_1. It contains no points of A_i. But $\tau_R^{-1}(M)$ is an A_i-module and so L is zero and g is a monomorphism. It is also an epimorphism. Thus a path in $\mathcal{X}_i$ is sectionally refinable in $\mathcal{X}_i$ if and only if it is also in ind$R$.

Moreover, let P be an A_i-modules in $\mathcal{X}_i$, if P is R-projective, then it is A_i-projective. If P is a R-projective module, than for every R-epimorphism, and in particular for every A_i-epimorphism $f : M \to N$ with $g : P \to N$, there exists an R-morphism $h : P \to M$ such that $hf = g$. Since h is a morphism between two A_i-modules, it is a morphism in modA_i. On the other hand, an A_i-module I in $\mathcal{X}_i$ is an injective R–module if and only if I is an injective A_i-module (this follows from Lemma 13 and from the construction of injective modules).

Now, let (Q, I) be a string bound quiver having a band and such that each band has only exiting arrows or only entering arrows. Let $R = kQ/I$ have no DOZE. We show that if M is not in $\mathcal{L}_R \cup \mathcal{R}_R$, then one of the following conditions is satisfied:

(1) The module M is in ind$C$;
(2) There exist i and j such that $1 \leq i \leq n$ and $1 \leq j \leq m$, and such that the module M is in indA_i∩indB_j;
(3) The module M is in indA_i, but not in $\mathcal{R}_{A_i}$;
(4) The module M is in indB_j, but not in $\mathcal{L}_{B_j}$.

If M does not satisfy the first two conditions, then by Proposition 20 there exists i such that M is in $\mathcal{X}_i$ or j such that M is in $\mathcal{Y}_j$. In the first case, if M is not in $\mathcal{R}_R$, then there exists a path of R-morphisms which is not sectionally refinable from M to an R-projective indecomposable module. From the results above, we obtain that M is not in $\mathcal{R}_{A_i}$. In the second, if M is not in $\mathcal{L}_R$, then there exists a path of R-morphisms which is not sectionally refinable from an R-injective indecomposable module to M. We obtain that M is not in $\mathcal{L}_{B_j}$. We have shown our statement.

In the four cases, M belongs to a finite set and thus all but a finite number of isomorphism classes of modules of ind$R$ are in $\mathcal{L}_R \cup \mathcal{R}_R$. In particular, R is laura. It cannot be quasi-tilted of canonical type since none of its bands have entering and exiting arrows (see Theorems 2.6 and 3.4 of [10]).
□

Corollary 21 *Let (Q, I) be a string bound quiver having bands and such that each band has only exiting arrows or only entering arrows. Then, if $R = kQ/I$ is a strict laura or a tilted string algebra without DOZE, its left and right end algebras are tilted of type $\widetilde{\mathbb{A}}_n$. Consequently, each string strict laura or tilted algebra without DOZE is domestic.*

Proof: We have that $\mathcal{X}_i$ is such that if M is in $\mathcal{X}_i$ and there exist a non-zero morphism from M to M', then M' is in $\mathcal{X}_i$, and such that only a finite number of A_i-modules are not in $\mathcal{X}_i$. Therefore, if R is not quasi-tilted, we have that $\mathcal{X}_i$ contains a complete slice of modA_i and this complete slice gives us a connected component of the right end algebra of R. Moreover, the right end algebra cannot contain another factor (in this case, by the arguments of the proof of Theorem 4, we obtain an infinite number of R-modules which are not in $\mathcal{L}_R \cup \mathcal{R}_R$). The domesticity of R follows from [2].
□

3 DOZED string modules

We now study string algebras whose bound quiver contains at least a DOZE.

Definition 22 *Let $R = kQ/I$ be a string algebra with a DOZE*

$$\alpha_1 ... \alpha_l \omega_1 {\omega_2}^n \omega_3 \beta_m ... \beta_1$$

for some walks $\alpha_1 \ldots \alpha_l$, ω_1, ω_3, $\beta_m ... \beta_1$ and a band ω_2. The string module $M_n = M(\sigma_n)$ corresponding to the string

$$\sigma_n = \alpha_3 ... \alpha_l \omega_1 {\omega_2}^n \omega_3 \beta_m ... \beta_3$$

is called the DOZED module of power n. By convention, $\alpha_3 ... \alpha_l$ represents the trivial path if the relation $\alpha_1 ... \alpha_l$ is of length 2.

Theorem 23 *Let R be a string algebra containing a DOZE. Then the DOZED string modules have projective and injective dimension greater than one. Thus, R is not laura.*

Proof: Let $\rho_1 \omega_1 \omega_2 \omega_3 \rho_2$ be a DOZE where ω_2 is a non-trivial band, $\rho_1 = \alpha_1 ... \alpha_l$ and $\rho_2 = \beta_m ... \beta_1$. Let $M_n = M(\sigma_n)$ be the DOZED module and I the injective

envelope of M_n.
Let $\alpha_3 ... \alpha_l \alpha_{l+1} ... \alpha_k$ be the longest path such that $\sigma_n = \alpha_3 ... \alpha_l \alpha_{l+1} ... \alpha_k \sigma'_n$ and x_3, x_4, ... x_{k+1} the vertices of this path. Then x_{k+1} is a sink of σ_n and thus $I_{x_{k+1}}$ is a direct summand of I. Let π_1 be the canonical projection on $I_{x_{k+1}}$, $M_n(x)$ and $M_n(\alpha)$ be the vector spaces and the linear maps corresponding to the vertex x and the arrow α in the representation M_n, respectively and f_{x_i} be the linear map associated to x_i in the representation of the morphism f. Then we have the following identities:

- $\pi_1 \circ M_n(\alpha_i) = \pi_{1 x_i}$ for every i such that $3 \leq i \leq k$;
- $\pi_1 \circ M_n(\alpha_2) = 0$ by definition of M_n;
- $\pi_1 \circ I(\alpha_i) = \pi_{1 x_i}$ for every i such that $2 \leq i \leq k$, since $\alpha_i ... \alpha_k$ is the first non-zero path going to $t(\alpha_k)$ which is a sink of σ_n;
- $\pi_1 \circ I(\alpha_1) = 0$ since $\alpha_1 ... \alpha_l$ is a relation ending in x_{l+1} and of minimal length for this property.

Let ι be the inclusion of M_n in I, ι_x the linear map induced by ι between $M_n(x)$ and $I(x)$, $C =$Cokerι and $c =$cokerι. Then:

- $\pi_1 \circ \iota_{x_i} = \pi_{1 x_i}$ for every i such that $3 \leq i \leq k$;
- $\pi_1 \circ \iota_{x_2} = 0$, by applying π_1 to the equation $\iota_{x_3} \circ M_n(\alpha_2) = I(\alpha_2) \circ \iota_{x_2}$;
- $\pi_1(C(x_2)) \cong k$ since $\pi_1 \circ \iota_{x_2} = 0$ and since $\pi_1(I(x_2)) \cong k$;
- $\pi_1 \circ c_{x_2} = \pi_{1 x_i}$;
- $\pi_1 \circ c_{x_3} = 0$;

$$\begin{tikzcd}
& \cdot \arrow[rr] \arrow[dd] & & \cdot \arrow[rr, "0"] \arrow[dd, "0"] & & \cdot \arrow[rr, "id_k"] \arrow[dd, "id_k"] & & \cdot \arrow[dd, "id_k"] \\
M_n(x_1) \arrow[ur, "\pi_1"] \arrow[rr, "M_n(\alpha_1)"] \arrow[dd, "\iota_{x_1}"] & & M_n(x_2) \arrow[ur, "\pi_1"] \arrow[rr, "M_n(\alpha_2)"] \arrow[dd, "\iota_{x_2}"] & & M_n(x_3) \arrow[ur, "\pi_1"] \arrow[rr, "M_n(\alpha_3)"] \arrow[dd, "\iota_{x_3}"] & & M_n(x_4) \arrow[ur, "\pi_1"] \arrow[dd, "\iota_{x_4}"] & \\
& \cdot \arrow[rr, "0"] \arrow[dd] & & \cdot \arrow[rr, "id_k"] \arrow[dd, "id_k"] & & \cdot \arrow[rr, "id_k"] \arrow[dd, "0"] & & \cdot \arrow[dd] \\
I(x_1) \arrow[ur, "\pi_1"] \arrow[rr, "I(\alpha_1)"] \arrow[dd, "c_{x_1}"] & & I(x_2) \arrow[ur, "\pi_1"] \arrow[rr, "I(\alpha_2)"] \arrow[dd, "c_{x_2}"] & & I(x_3) \arrow[ur, "\pi_1"] \arrow[rr, "I(\alpha_3)"] \arrow[dd, "c_{x_3}"] & & I(x_4) \arrow[ur, "\pi_1"] \arrow[dd, "c_{x_4}"] & \\
& \cdot \arrow[rr, "0"] & & k \arrow[rr, "0"] & & \cdot \arrow[rr] & & \cdot \\
C(x_1) \arrow[ur, "\pi_1"] \arrow[rr, "C(\alpha_1)"] & & C(x_2) \arrow[ur, "\pi_1"] \arrow[rr, "C(\alpha_2)"] & & C(x_3) \arrow[ur, "\pi_1"] \arrow[rr, "C(\alpha_3)"] & & C(x_4) \arrow[ur, "\pi_1"] &
\end{tikzcd}$$

By applying π_1 to the equation $C(\alpha_1) \circ c_{x_1} = c_{x_2} \circ I(\alpha_1)$, we obtain that $\pi_1 \circ C(\alpha_1) \circ c_{x_1} = \pi_1 \circ c_{x_2} \circ I(\alpha_1) = \pi_1 \circ I(\alpha_2) = 0$. Since c_{x_1} is an epimorphism, we have $\pi_1 \circ C(\alpha_1) = 0$. We show in the same way that $\pi_1 \circ C(\alpha_2) = 0$, and that $\pi_1 \circ C(\beta) = 0$ for every arrow β having x_2 for its source.
Thus C admits an indecomposable direct summand whose support does not contain x_1 and admits x_2 as a sink. So this direct summand (and thus C) is

not an injective module.
□

Proof of Theorem 5: a) implies b):

(1) If there is no band, the algebra is of finite representation type by [6], and we are done;
(2) Otherwise, Theorem 4 gives us the statement.

The statement b) implies trivially c). The statement c) implies a) by Theorem 23.
□

We now generalize Theorem 5 to special biserial algebras.

4 Laura special biserial algebras and Skowroński's conjecture

Definition 24 *Let R be an algebra. It is a special biserial algebra if it admits a presentation $R = kQ/I$ such that :*

(1) Each point has at most two arrows entering and two arrows exiting;
(2) For each arrow $\alpha : x \to y$ there is at most one arrow $\beta : y \to z$ such that $\alpha\beta$ is not in I and at most one arrow $\gamma : z \to x$ such that $\gamma\alpha$ is not in I.

Remark: For a special biserial algebra R, we denote J the ideal generated by the paths appearing in the commutativity relations of I. Then, R/J is a string algebra.

Definition 25 *Let $R = kQ/I$ be a special biserial algebra and (ρ_1, ρ_2) be two consecutive zero-relations on Q. We say that they form a DOZE if they do so on R/J.*

By [13], if R is special biserial, any indecomposable R-module which is not projective-injective is also an R/J-module. This implies that any indecomposable R-module which is not projective-injective is a string module or a band module. Moreover, the restriction of scalars functor $\text{mod}R/J \to \text{mod}R$ is full, faithful and sends an irreducible morphism in $\text{mod}R/J$ onto an irreducible morphism in $\text{mod}R$.

Proof of Theorem 6: The proof that (b) implies (c) follows from the definition of laura algebras.
Suppose that there exists a DOZE on (Q, I) and let M_n be a DOZED module on R/J. Then M_n is also an R-module. We have shown in the proof of Theorem 23 that the injective dimension over R/J of M_n is greater than or equal to

two by building $I_{x_{l+1}}$ and the cokernel C. Remark that a DOZE on R contains no path involved in a binomial relation. By applying the same technique, we show that the injective dimension of M_n over R is greater than or equal to two and we get that c) implies a).
The proof that a) implies d) follows from Theorem 23.
For the last implication, let R be a special biserial algebra such that R/J is laura. Let M be a non projective-injective R-module which is not in $\mathcal{L}_R \cup \mathcal{R}_R$. Then there exist a non sectionally refinable path (*) going from M to a R-projective module P and a non sectionally refinable path (**) going from an R-injective module I to M.

$$\begin{tikzcd}[column sep=small]
(*) & M \arrow[dr] & & \tau^{-1}M \arrow[r, squiggly] & P & & (**) & I \arrow[r, squiggly] & \tau M \arrow[dr] & & M \\
& & \cdot \arrow[ur] & & & & & & & \cdot \arrow[ur] &
\end{tikzcd}$$

The fact that (*) and (**) are not sectionally refinable is preserved in R/J since an almost split sequence admitting a projective-injective middle term has at least another middle term [4]. Moreover, if P and I are not projective-injective modules, then they are R/J-projective and R/J-injective modules respectively. If P is projective-injective, then $P/\mathrm{soc}P$ is an R/J-projective module. We only have to compose (*) with the left minimal almost split morphism going from P to $P/\mathrm{soc}P$, and we get the desired path from M to an R/J-projective module.

$$\begin{tikzcd}
M \arrow[r, squiggly] & P \arrow[r] & P/socP
\end{tikzcd}$$

In all cases, if M is a non projective-injective R-module which is not in $\mathcal{L}_R \cup \mathcal{R}_R$, then it is not in $\mathcal{L}_{R/J} \cup \mathcal{R}_{R/J}$, and so the set of indecomposable R-modules which are not in $\mathcal{L}_R \cup \mathcal{R}_R$ is finite.
□

We conclude with an example.

Example: Let $R = kQ/I$, where Q is the following quiver:



and I is generated by $\alpha_i\beta_i$ and $\gamma_i\delta_i$. This algebra is not a string algebra, but by applying the evident action of the group $\mathbb{Z}_2$, we obtain the skew group algebra $A[G]$ given by the following quiver:

$$\begin{tikzcd}
 & & x_4 \arrow[dr, "\gamma_1"] \arrow[drr, dash, dotted, bend left] & & \\
x_1 \arrow[r, "\alpha"] \arrow[urr, dash, dotted, bend left] \arrow[drr, dash, dotted, bend right] & x_2 \arrow[ur, "\beta_1"] \arrow[dr, "\beta_2"'] & & x_5 \arrow[r, "\delta"] & x_6 \\
 & & x_3 \arrow[ur, "\gamma_2"'] \arrow[urr, dash, dotted, bend right] & &
\end{tikzcd}$$

where I is generated by $\alpha\beta_1$, $\alpha\beta_2$, $\gamma_2\delta$ and $\gamma_1\delta$. It is a string algebra which contains the DOZE $\alpha\beta_1\gamma_1\delta$, this latter algebra is not laura by Theorem 5 and so the initial algebra R is not laura by [3].

ACKNOWLEDGEMENTS. The author warmly thanks Ibrahim Assem as well as Patrick Le Meur and Shiping Liu for their suggestions. She also thanks ISM and Sherbrooke University for their financial support.

References

[1] I. Assem and F. Coelho. Glueings of tilted algebras. *J. Pure and Appl. Algebra*, 96(3):225–243, 1994.

[2] I. Assem and F. Coelho. Two-sided gluings of tilted algebras. *J. Algebra*, 269(2):456–479, 2003.

[3] I. Assem, M. Lanzilotta, and M.J. Redondo. Laura skew group algebras. *Comm. Algebra*, 35(7):2241–2257, 2007.

[4] I. Assem, D. Simson, and A. Skowroński. *Elements of Representation Theory of Associative Algebras.* London Math. Soc. Students Texts. Cambridge University Press.

[5] J. Bélanger and C. Tosar. Shod string algebras. *Comm. Algebra*, 33(8):2465–2487, 2005.

[6] M.C.R. Butler and C.M. Ringel. Auslander-Reiten sequences with few middle terms ; Applications to string algebras. *Comm. Algebra*, 15(1-2):145–179, 1987.

[7] K. Fuller. Biserial rings. *Lecture Notes in Math.*, (734):64–90, 1979.

[8] D. Happel. A characterization of hereditary categories with tilting object. *Invent. Math.*, 144(2):381–398, 2001.

[9] F. Huard and S. Liu. Tilted Special Biserial Algebras. *J. Algebra*, 217(2):679–700, 1999.

[10] F. Huard and S. Liu. Tilted String Algebras. *J. Pure Appl. Algebra*, 153:151–164, 2000.

[11] A. Skowroński. On Artin algebras with almost all indecomposable modules of projective or injective dimension at most one. *Cent. Eur. J. Math.*, 1(1):108–122, 2003.

[12] A. Skowroński and J. Waschbüsch. Representation-finite biserial algebras. *J. reine angew. Math.*, (345):172–181, 1983.

[13] B. Wald and I. Waschbüsch. Tame biserial algebras. *J. Algebra*, 95:480–500, 1985.